



HIPPOKRATIA
Quarterly Medical Journal
ISSN 1108-4189



Note: Performing your original search, diabetes leading cause acute renal failure, in PubMed 223 citations.

Journal List > Hippokratia > v.12(1); Jan-Mar 2008

Abstract

Full Text

PDF (879K)

Contents

Archive

Related material:

PubMed related arts

GO

PubMed articles by:
Sampanis, C.

Top

Abstract

The problem of
diabetic nephropathy
The role of kidney in
the metabolism of
insulin in normal
man and in renal
failure

The role of insulin
resistance

Insulin secretion in
renal failure

Insulin requirements
in diabetes mellitus
with renal disease

Glycemic control in
patients with
diabetes mellitus and
renal failure

Management of
patients with
diabetes mellitus and
advanced kidney
disease

Pharmacological
treatment of

Hippokratia, 2008 Jan-Mar; 12(1): 22-27.

Copyright 2008, Hippokratia General Hospital of Thessaloniki

Management of hyperglycemia in patients with diabetes mellitus and chronic renal failure

Ch Sampanis

Diabetes Center, 2nd Department of Internal Medicine, Aristotle University of Thessaloniki, H
Hospital,

Sampanis Ch, 9 Glinou Str, 54249, Thessaloniki, Greece, Tel.+302310311928

Abstract

Diabetes mellitus is recognized as a leading cause of chronic kidney disease and failure. Chronic renal failure is associated with insulin resistance and, in advanced stages, decreased insulin degradation. Both of these abnormalities are partially reversed by dialysis. Except for diet with protein restriction, patients with diabetes should be treated with insulin. The management of the patients with hyperglycemia and chronic renal failure requires close collaboration between the diabetologist and the nephrologists. This collaboration is important so that the patient will not be confused and will not lose confidence in the medical team. Furthermore good glycemic control in these patients seems to reduce microvascular complications.

Keywords: hyperglycemia therapy, diabetes, chronic kidney disease

hyperglycemia in
patients with
diabetes mellitus and
advanced kidney
disease

Use of insulin in
chronic renal failure

References

Top

Abstract

The problem of
diabetic nephropathy
The role of kidney in
the metabolism of
insulin in normal
man and in renal
failure

The role of insulin
resistance

Insulin secretion in
renal failure

Insulin requirements
in diabetes mellitus
with renal disease

Glycemic control in
patients with
diabetes mellitus and
renal failure

Management of
patients with
diabetes mellitus and
advanced kidney
disease

Pharmacological
treatment of
hyperglycemia in
patients with
diabetes mellitus and
advanced kidney
disease

Use of insulin in
chronic renal failure

References

Top

Abstract

■ The problem of

Diabetes mellitus is a major health problem of increasing magnitude worldwide on cardiovascular morbidity and mortality¹. Moreover diabetes mellitus is recognized as a cause of chronic kidney disease and end-stage renal disease in the United States and other Countries². Large epidemiological studies have shown that one third of the patients with end-stage renal transplant recipients are diabetics, predominantly with type 2 diabetes³. Large designed randomized studies have provided convincing evidence on the value of preventing both micro and macrovascular disease⁴. The UK Prospective Diabetes Study (UKPDS) that intensive treatment of patients with newly diagnosed diabetes reduced the risk of myocardial infarction by 16%, amputation or death from peripheral vascular disease by 35%, stroke by 42%, nonfatal myocardial infarction by 21%, fatal sudden death by 39%. Every 1% reduction in glycosylated hemoglobin was associated with a 12% reduction in risk of myocardial infarction and 14% for diabetes related deaths, 14% for microvascular complications. Therefore the glycemic control is an important factor for the prevention of diabetic complications^{5,6}.

The problem of diabetic nephropathy

In the past it has been believed that fewer patients with type 2 diabetes develop

diabetic nephropathy

The role of kidney in the metabolism of insulin in normal man and in renal failure

The role of insulin resistance

Insulin secretion in renal failure

Insulin requirements in diabetes mellitus with renal disease

Glycemic control in patients with diabetes mellitus and renal failure

Management of patients with diabetes mellitus and advanced kidney disease

Pharmacological treatment of hyperglycemia in patients with diabetes mellitus and advanced kidney disease

Use of insulin in chronic renal failure

References

Top

Abstract

The problem of

diabetic nephropathy

■ The role of kidney in the metabolism of insulin in normal man and in renal failure

The role of insulin resistance

Insulin secretion in renal failure

Insulin requirements in diabetes mellitus with renal disease

that proteinuria in these patients had relatively better prognosis compared to pa diabetes. Well designed prospective studies have shown that once proteinuria d end-stage renal disease is similar in both types of diabetes³. Moreover recent ep have shown that end-stage renal failure has increased dramatically in patients w The reason for this is that the treatment of hypertension and coronary heart dise life expectancy of the patients with type 2 diabetes and larger proportion of the nephropathy and end-stage renal disease^{4,9}.

The role of kidney in the metabolism of insulin in and in renal failure

In non-diabetic individual, 40-50% of insulin secreted by pancreas is extracted passage through the liver^{10,11}. Consequently, the kidney plays a smaller role in c secreted in non-diabetic individual than in disposing of insulin injected into dia 1). Endogenously secreted insulin is degraded by liver, exogenous insulin is pri the kidney.




Figure 1.

Metabolism of insulin. Insulin is freed at the glomerulus and then e reabsorbed by the proximal tubule. Of the total renal insulin clearanc 60% occurs by glomerular filtration and 40% by extraction from peri

Glycemic control in
patients with
diabetes mellitus and
renal failure

Abstract
Management of
patients with
diabetes mellitus and
advanced kidney
disease

Pharmacological
treatment of
hyperglycemia in
patients with
diabetes mellitus and
advanced kidney
disease

Use of insulin in
chronic renal failure

References

(more ...)

The kidneys play an important role in the clearance of insulin from the systemic has a molecular weight of 5734 and is therefore freely filtered at the glomerulus reabsorbed by the proximal tubule. Of the total renal insulin clearance, approxir glomerular filtration and 40% by extraction from peritubular vessels. Insulin in enters the proximal tubular cell by carrier-mediated endocytosis and is then tran lysosomes where it is metabolized into amino acids that are released into peritil diffusion. In addition to luminal clearance via glomerular filtration, the kidneys post-glomerular peritubular circulation (Figure 2). These intrarenal pathways of involve both receptor and non-receptor mediated uptake. The net effect is that l insulin appears in final urine¹²⁻¹⁴.



Figure 2.

Intrarenal pathways of insulin removal. Filtered insulin is internalized endocytosis and thereafter degraded into amino acids into the peritub Insulin removed from postglomerular peritubular vessels binds to the cell membrane. (more ...)

In patients with advanced renal failure basal plasma levels of insulin, proinsulin elevated. The renal clearance of C-peptide is greater than of insulin in renal inst the use of C-peptide concentration as an index of insulin secretion in these patie During the early phase of renal insufficiency, impaired renal insulin clearance is blood flow, but as renal function declines, the effect of reduced blood flow is ag of tissue extraction of insulin¹³. As renal failure progresses, peritubular insulin i compensates for the decline in degradation of filtered insulin until the glomerul decreases to less than about 20 ml/min, after which insulin clearance decreases, increases, and overall requirements for insulin decline^{12,16-18}. There is evidence that renal failure suppresses insulin metabolism in extrarenal sites, e.g. in skelet liver. Insulin-dependent diabetic patients often occur a decrease in insulin requi patients with residual β cell function, the need for exogenous insulin may disap with endstage renal disease and hemodialysis occur hypoglycemia because of p of circulating insulin, altered dietary and exercise patterns²⁰.

Under normal circumstances, the capacity of renal tubules to absorb filtered insi saturation does not occur, therefore insulin clearance normally is constant over i concentration.

The role of insulin resistance

Patients with renal failure have impaired insulin sensitivity with consequent abn metabolism. The underlying mechanism is not clear, but an increase of glucone

Top

Abstract

The problem of
diabetic nephropathy

The role of kidney in the metabolism of insulin in normal man and in renal failure

■ The role of insulin resistance

Insulin secretion in renal failure

Insulin requirements in diabetes mellitus with renal disease

Glycemic control in patients with diabetes mellitus and renal failure

Management of patients with diabetes mellitus and advanced kidney disease

Pharmacological treatment of hyperglycemia in patients with diabetes mellitus and advanced kidney disease

Use of insulin in chronic renal failure

References

Top

Abstract

The problem of diabetic nephropathy

The role of kidney in the metabolism of insulin in normal man and in renal failure

The role of insulin resistance

■ Insulin secretion in renal failure

Insulin requirements in diabetes mellitus with renal disease

reduction of hepatic and/or skeletal muscle glucose uptake and an impairment of glucose metabolism due either to decreased oxidation to carbon dioxide and water. The synthesis of glycogen may be involved^{18,21,22}. The exact mechanism of insulin resistance with renal failure is still debated. There are some controversies in the literature, clinical studies indicate that in uremia glucose production and uptake in the liver and skeletal muscle is the primary site of insulin resistance^{18,23}. Moreover, glucocorticoid abnormality seems to be of great importance, as the rate of glucose oxidation is

Other factors contributing to insulin resistance in uremic patients are an accumulation of toxins and an excess of parathyroid hormone. The role of uremic toxins was suggested by the observation that hemodialysis improved insulin sensitivity²⁴⁻²⁶. Some studies have shown that intravenous therapy with calcitriol (1,25-dihydroxyvitamin D) and hyperparathyroidectomy enhanced insulin sensitivity and improved glucose tolerance. Decreased tissue oxygenation because of anemia also contributes to insulin resistance in uremia as evidenced by the increase in insulin sensitivity after correction of anemia by erythropoietin^{27,28}.

Insulin secretion in renal failure

As glomerular filtration decreases below 50 ml/min insulin secretion seems to be probably due to the presence of metabolic acidosis, the parathyroid hormone exerts the elevation of intracellular calcium concentration and decrease of the cellular Na-K-ATPase pump activity in the pancreatic β cells. Experimental studies have shown that changes may be prevented by prior parathyroidectomy or by the administration of the channel blocker verapamil and that the effect of metabolic acidosis on insulin secretion is reversed by hemodialysis^{23,29-31}.

There are some controversies about the effect of erythropoietin therapy on insulin secretion. Some studies showed an increase of insulin secretion and decrease of blood glucose levels, while other studies showed no change in insulin secretion after an oral glucose load.

Glycemic control in patients with diabetes mellitus and renal failure

Management of patients with diabetes mellitus and advanced kidney disease

Pharmacological treatment of hyperglycemia in patients with diabetes mellitus and advanced kidney disease

Use of insulin in chronic renal failure

References

Top

Abstract

The problem of diabetic nephropathy

The role of kidney in the metabolism of insulin in normal man and in renal failure

The role of insulin resistance
Insulin secretion in renal failure

■ Insulin requirements in diabetes mellitus with renal disease

Glycemic control in patients with diabetes mellitus and renal failure

Management of patients with diabetes mellitus and advanced kidney disease

Pharmacological treatment of hyperglycemia in

Insulin requirements in diabetes mellitus with re

Insulin requirements show a biphasic course in patients with diabetes and renal beginning glucose control deteriorates because of insulin resistance, therefore n to achieve glycemic control. In advanced renal failure with creatinine clearance need for insulin is lower or even the cessation of insulin may be necessary. The decreased because of less caloric intake in uremic patients. With the institution need for insulin changes because the insulin sensitivity and liver metabolism in

In both non-diabetic and diabetic subjects with chronic renal failure spontaneous develop because of decreased caloric intake, reduced renal gluconeogenesis, im counterregulatory hormone epinephrine due to the autonomic neuropathy, conc and decreased metabolism of drugs that might promote a reduction in plasma gl such as alcohol, nonselective blockers, and disopyramide^{23,36,37}. The plasma eli significantly increased in patients with renal insufficiency when clearance of cr 40ml/min, therefore in these patients the dose adjustment of disopyramide is ne

patients with
diabetes mellitus and
advanced kidney
disease

Use of insulin in
chronic renal failure

References

Top

Abstract

The problem of
diabetic nephropathy

The role of kidney in
the metabolism of
insulin in normal
man and in renal
failure

The role of insulin
resistance

Insulin secretion in
renal failure

Insulin requirements
in diabetes mellitus
with renal disease

■ Glycemic control in
patients with diabetes
mellitus and renal
failure

Management of
patients with
diabetes mellitus and
advanced kidney
disease

Pharmacological
treatment of
hyperglycemia in
patients with
diabetes mellitus and
advanced kidney
disease

Use of insulin in
chronic renal failure

References

Top

Abstract

The problem of
diabetic nephropathy

Glycemic control in patients with diabetes mellitus failure

Measurement of the glycated hemoglobin (HbA1c) seems to be the most accurate measure of glycemic control in patients with diabetes. However there are some limitations to its use, due to interference from carbamylated hemoglobin that leads to a falsely low HbA1c level. Other factors that affect the accuracy of the HbA1c measurement include recent transfusion, iron deficiency, accelerated erythropoiesis due to therapy, and metabolic acidosis³⁸⁻⁴³. Despite these limitations, results on the range of HbA1c values appear to estimate glycemic control similarly to patients without advanced renal failure, while values over 7.5% may overestimate the extent of hyperglycemia.

Self blood glucose monitoring (SMBG) permits estimation of chronic glycemic control. The American Diabetes Association (ADA) recommends that patients with type 1 diabetes monitor blood glucose at least three times daily, and type 2 diabetes patients that are treated with hypoglycemic drugs monitor blood glucose daily^{44,45}. Self-blood glucose monitoring is important in type 1 diabetes, because their blood glucose concentrations are less stable than those in patients with type 2 diabetes.

Fasting blood glucose levels < 140 mg/dl, < 200 mg/dl one hour after meal and between 6-7% in type 1 diabetes and between 7-8% in patients with type 2 diabetes are considered acceptable⁴⁶.

Excellent glycemic control has not been emphasized as much in diabetic dialysis patients without renal failure because of the possible precipitation of hypoglycemia, especially with fluctuating dietary intake⁴⁷.

Management of patients with diabetes mellitus and kidney disease

The management of diabetic patients with advanced kidney disease includes diet

The role of kidney in the metabolism of insulin in normal man and in renal failure

The role of insulin resistance

Insulin secretion in renal failure

Insulin requirements in diabetes mellitus with renal disease

Glycemic control in patients with diabetes mellitus and renal failure

■ Management of patients with diabetes mellitus and advanced kidney disease

Pharmacological treatment of hyperglycemia in patients with diabetes mellitus and advanced kidney disease

Use of insulin in chronic renal failure

References

Top

Abstract

The problem of diabetic nephropathy

The role of kidney in the metabolism of insulin in normal man and in renal failure

The role of insulin resistance

Insulin secretion in renal failure

Insulin requirements in diabetes mellitus with renal disease

Glycemic control in

restriction and oral agents or insulin. Patients with renal failure are often confused by recommendations and have the impression that different specialists have contradictory advice.

In patients with type 1 diabetes insulin therapy imposes regularity in food intake and intake of carbohydrates. A dietary pattern in these patients should include carbohydrates, vegetables, whole grains, legumes, and low-fat milk. Moreover, monitoring carbohydrate counting, exchanges or experienced-based estimation remains a key to achieving glycemic control and maintain a stable body weight.

Most of the patients with type 2 diabetes are obese, with insulin resistance and hypersecretion. These patients must be encouraged to lose weight with hypocaloric diet. Protein restriction is in practice replaced by carbohydrates or fat in order to maintain caloric intake. It has been shown that both short-term and long-term increases in protein ratio are accompanied by an improvement in the action of insulin⁴⁹. However, when the ratio of carbohydrates to protein is increased, hypertriglyceridemia may develop. On the other hand, increasing the ratio of saturated fat to protein was accompanied by the inhibition of insulin action, while the ratio of monounsaturated or polyunsaturated fats did not rise the insulin resistance⁵⁰. To appropriately compensate this reduction in protein caloric intake by carbohydrates of both types of diabetes with renal failure should be advised to go on a protein-restricted diet to compensate the loss in calories by carbohydrates. The dietary management in type 2 diabetes requires close collaboration between the dietician, the diabetologist and the nephrologist.

Pharmacological treatment of hyperglycemia in patients with diabetes mellitus and advanced kidney disease

The antihyperglycemic drugs available for the treatment of type 2 diabetes include sulphonylureas and meglitinides, metformin, α -glucosidase inhibitors, the thiazolidinediones and insulin, while patients with type 1 diabetes mellitus are treated with insulin.

Agent	Half-life (h)	Protein binding (%)	Renal clearance (ml/min)
Glibenclamide	6-12	99	10-20
Gliclazide	12-18	99	10-20
Glimipride	12-18	99	10-20
Glibenclamide	6-12	99	10-20
Gliclazide	12-18	99	10-20
Glimipride	12-18	99	10-20
Glibenclamide	6-12	99	10-20
Gliclazide	12-18	99	10-20
Glimipride	12-18	99	10-20

Table 1

Oral anti hyperglycemic agents (adapted from RW Snyder and JS Berkenstadt, *Diabetes* 2004, Vol 17, No 5, pp 365-370)

Sulphonylureas (Glyburide, Glipizide, Glimipride) are strongly protein bound, therefore elevated drug plasma levels cannot be efficiently reversed by dialysis.

patients with
diabetes mellitus and
renal failure

Management of
patients with
diabetes mellitus and
advanced kidney
disease

■ Pharmacological
treatment of
hyperglycemia in
patients with diabetes
mellitus and advanced
kidney disease

Use of insulin in
chronic renal failure

References

administration of these drugs in patients with end-stage renal disease requires caution and the routes of elimination.⁵¹

Glyburide has weak metabolites that are excreted in the urine and accumulate in renal failure. This drug can be given at reduced dose if the GFR is above 50 ml/min, in more severe renal impairment.

Glipizide is metabolized in the liver into several inactive metabolites. Less than 1% of Glipizide is excreted unchanged in urine and about 60% is excreted as metabolites. Metabolites accumulate in patients with renal failure and, one of them may have activity, that does not cause hypoglycemia. Moreover, Glipizide clearance and half-life (two to four hours), therefore dose adjustment in patients with reduced GFR. Glipizide should probably be the sulphonylurea of choice in patients with renal failure as it is not available in Greek market at present time.

Glimepiride is metabolized in the liver, the elimination half-time is about 5-8 hours. A small amount of a dose appears in the urine. Virtually all of the urinary excretion is as metabolites. In patients with renal failure. Glimepiride may cause prolonged hypoglycemia in patients with renal dysfunction, so the drug should be used cautiously, and the dose of this drug must be reduced in patients with a reduced GFR.

Repaglinide is a drug that is exclusively metabolized in the liver, with a half-life of about 4 hours and is excreted in the bile and stool. Only 10% of the total dose appears in the urine. Concentration and elimination half-life are increased in patients with renal failure. Nevertheless, dose reduction is not necessary in these patients⁵⁴.

Nateglinide, is a drug with hepatic metabolism and elimination half-life of about 2 hours. Several of the metabolites are active and accumulate and in patients with impaired renal function causing hypoglycemia. Therefore this drug should be used cautiously in such patients.

Metformin is primarily excreted unchanged in the urine. Patients with renal failure are susceptible to drug accumulation and lactic acidosis. Therefore this drug is contraindicated if creatinine clearance is below 60 ml/min⁵¹.

The thiazolidinediones (Pioglitazone, Rosiglitazone) are virtually completely metabolized in the liver, each forming several metabolites. For both drugs there is no accumulation of the major metabolites in the setting of renal insufficiency. These drugs may cause congestive heart failure. Plasma volume increases in patients treated with thiazolidinediones consequent anemia due to hemodilution. The risk for edema is more prominent when used in patients in combination with insulin^{51,57,58}. Given the risk of edema formation, these drugs should be avoided in patients with advanced kidney disease, have preexisting heart failure. Whether these drugs should be avoided in patients with renal failure is uncertain.

Alpha-glucosidase inhibitors such as acarbose or miglitol are renally excreted and accumulate in renal dysfunction and are contraindicated in patients with renal

Use of insulin in chronic renal failure

Top

Abstract

The problem of
diabetic nephropathy

The role of kidney in
the metabolism of
insulin in normal
man and in renal
failure

The role of insulin
resistance

Insulin secretion in
renal failure

Insulin requirements
in diabetes mellitus
with renal disease

Glycemic control in
patients with
diabetes mellitus and
renal failure

Management of
patients with
diabetes mellitus and
advanced kidney
disease

Pharmacological
treatment of
hyperglycemia in
patients with
diabetes mellitus and
advanced kidney
disease

■ Use of insulin in
chronic renal failure

References

Chronic renal failure is associated with decreased renal and hepatic metabolism decreased clearance and metabolism of insulin, the metabolic effects of insulin longer and the risk for hypoglycemia increases. According to current recommen adjustment is required if the GFR is above 50 ml/min. The insulin dose should be approximately 75% when the GFR is between 10-50 ml/min and by as much as is less than 10 ml/min^{51,52,59}.

It is of great importance that the net effect on glycemic control will vary from patient to patient therefore dose adjustment may be necessary.

The 2005 K/DOQI guidelines suggest that in patients with end-stage renal disease should be encouraged for glycemic control, and oral antihyperglycemic drugs with exception of Glipizide⁶⁰.

Insulin regimens are the same as in patients without renal insufficiency^{51,61,62}.

The pharmacokinetics of various insulin preparations have not been well studied in varying degrees of renal dysfunction, and there are no absolute guidelines defining dosing adjustment of insulin that should be made based on the level of GFR⁵¹. In stage renal disease some suggest that long-acting insulin preparations should be used with support that such agents should be used.

Patients treated with continuous ambulatory peritoneal dialysis or continuous cyclic peritoneal dialysis (CAPD and CCPD) can be treated with intraperitoneal insulin. This regimen of continuous insulin infusion, eliminates the need for injections and provides a means of absorption⁶³⁻⁶⁵. The disadvantage of this therapy is that there is an additional contamination of dialysate during injection of the insulin into the bags, the need for insulin because of losses of spent dialysate and increased risk of peritoneal proliferation⁶⁶ and hepatic subcapsular steatosis⁶³. Moreover the absorption of insulin significantly varies among patients or may decline over time due to acquired abnormalities of the peritoneal membrane.

In conclusion patients with diabetes mellitus and advanced kidney disease should be treated with protein restriction and, preferably by insulin regimen. Oral hypoglycemic agents should be avoided because of risk of hypoglycemia, with exception of glipizide or repaglinide should be adjusted on the basis of glucose values obtained by home glucose monitoring. In addition to self-monitoring of blood glucose, while performing glycated hemoglobin (HbA1c) permits estimation of chronic glycemic control.

Top

Abstract

The problem of
diabetic nephropathy

The role of kidney in
the metabolism of
insulin in normal
man and in renal
failure

The role of insulin
resistance

Insulin secretion in
renal failure

Insulin requirements
in diabetes mellitus
with renal disease

Glycemic control in
patients with
diabetes mellitus and
renal failure

Management of
patients with
diabetes mellitus and
advanced kidney
disease

Pharmacological
treatment of
hyperglycemia in
patients with
diabetes mellitus and
advanced kidney
disease

Use of insulin in
chronic renal failure

■ References

References

1. Grundy SM, Howard B, Smith SJr, Eckel R, Redberg R, Bonow RO. Prevention Confer Cardiovascular Disease: executive summary: conference proceeding for healthcare prof writing group of the American Heart Association. *Circulation*. 2002;105:2231-2239. [P]
2. Mokdad AH, Serdula MK, Dietz WH, et al. The spread of the obesity epidemic in the U
JAMA. 1999;282:1519-1522. [PubMed]
3. *USRDS 1999 annual data report*. Bethesda, Md: National Institutes of Health, National Digestive and Kidney Disease; 1999. US Renal Data System; pp. 25-38.
4. Gerstein HC. Dysglycaemia: a cardiovascular risk factor. *Diabetes Res Clin Pract*. 1998 [PubMed]
5. UK Prospective Diabetes Study Group. Intensive blood-glucose control with sulphonylh with conventional treatment and risk of complications in patients with type 2 diabetes. (1998;352:837-853. [PubMed]
6. Stratton IM, Adler AI, Neil AW, et al. Association of glycaemia with microvascular anc complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ*. 20 [PubMed]
7. Canadian Diabetes Association Clinical Practice Guidelines Expert Committee. Canad 2003 Clinical Practice Guidelines for the Prevention and Management of Diabetes in Ca 2003;27(Suppl 2):s21-s23.
8. Hasslacher C, Ritz E, Wahl P, Michael C. Similar risks of nephropathy in patients with 1 mellitus. *Nephrol Dial Transplant*. 1989;4:859-863. [PubMed]
9. Ritz E, Stefanski A. Diabetic nephropathy in type II diabetes. *Am J Kidney Dis*. 1996;27
10. Rabkin R, Ryan MP, Duckworth WC. The renal metabolism of insulin. *Diabetologia*. 1⁴
11. Ferrannini E, Wahren J, Faber OK, Felig P, Binder C, DeFronzo RA. Splanchnic and re in human subjects: a dose-response study. *Am J Physiol*. 1983;244:E517-E527. [PubMed]
12. Rubenstein AH, Mako ME, Horowitz DL. Insulin and the kidney. *Nephron*. 1975;15:30
13. Rabkin R, Simon NM, Steiner S, Colwell JA. Effect of renal disease on renal uptake and man. *N Engl J Med*. 1970;282:182-187. [PubMed]
14. Katz AI, Rubenstein AH. Metabolism of proinsulin, insulin and C-peptide in rat. *J Clin* 1121. [PubMed]
15. Jaspén JB, Mako ME, Kuzuya H, Blix PM, Horowitz DL, Rubenstein AH. Abnormalitie peptides in chronic renal failure: comparison of C-peptide, proinsulin and insulin. *J Clin*

1977;45:441–446. [PubMed]

16. Rave K, Heise T, Pflutzner A, Heinemann L, Sawicki PT. Impact of diabetic nephropathy pharmacokinetic properties of insulin in type 1 diabetic patients. *Diabetes Care*. 2001;24:1033–1038. [PubMed]
17. Biesenbach G, Raml A, Schmckal B, Eichbauer-Sturm G. Decreased insulin requirement in nephropathic type 1 and insulin-treated type 2 diabetic patients. *Diabet Med*. 2003;20:641–646. [PubMed]
18. Mak RH, DeFronzo RA. Glucose and insulin metabolism in uremia. *Nephron*. 1992;61:1–10. [PubMed]
19. Maack T, Johnson V, Kan ST, Figueiredo J, Sigulem D. Renal filtration, transport and molecular weight proteins: a review. *Kidney Int*. 1979;16:251–270. [PubMed]
20. Goldberg AP, Hagberg JM, Delmez JA, Haynes ME, Harter HR. Metabolic effects of continuous hemodialysis patients. *Kidney Int*. 1980;18:754–761. [PubMed]
21. Alvestrand A. Carbohydrate and insulin metabolism in renal failure. *Kidney Int*. 1997;6:1–10. [PubMed]
22. Carone FA, Peterson DR. Hydrolysis and transport of small peptides by the proximal tubule. *Kidney Int*. 1980;238:F151–158. [PubMed]
23. Adrogu HJ. Glucose homeostasis and the kidney. *Kidney Int*. 1992;42:1266–1272. [PubMed]
24. Mak RH. Intravenous 1,25-dihydroxycholecalciferol corrects glucose intolerance in hemodialysis patients. *Kidney Int*. 1992;41:1049–1054. [PubMed]
25. Kautzsky-Willer A, Pacini G, Barnas U, et al. Intravenous calcitriol normalizes insulin sensitivity in hemodialysis patients. *Kidney Int*. 1995;47:200–206. [PubMed]
26. Lin S, Lin Y, Lu K, et al. Effects of intravenous calcitriol on lipid profiles and glucose tolerance in patients with secondary hyperparathyroidism. *Clin Sci*. 1994;87:533–538. [PubMed]
27. Borissova AM, Djambazova A, Todorov K, Dakovska L, Tankova T, Kirilov G. Effect of continuous hemodialysis on metabolic state and peripheral insulin sensitivity in diabetic patients on hemodialysis. *Nephrol Dial Transplant*. 1993;8:93–98. [PubMed]
28. Mak RH. Effect of recombinant human erythropoietin on insulin, amino acid, and lipid metabolism in hemodialysis patients. *Pediatr*. 1996;129:97–104. [PubMed]
29. Fadda GZ, Hajjar SM, Perna AF, Zhou XJ, Lipson LG, Massry SG. On the mechanism of insulin secretion in chronic renal failure. *J Clin Invest*. 1991;87:255–261. [PubMed]
30. Perna AF, Fadda GZ, Zhou XJ, Massry SG. Mechanisms of impaired insulin secretion in chronic renal failure. *Am J Physiol*. 1990;259:F210–206. [PubMed]
31. Oh H, Fadda GZ, Smogorzewski M, Liou HH, Massry SG. Abnormal leucine-induced insulin secretion in chronic renal failure. *Am J Physiol*. 1994;267:F853–860. [PubMed]
32. Kokot F, Wiecek A, Grzeszczak W, Klin M, Zukovska-Szczecawska E. Influence of continuous hemodialysis on glucose tolerance, insulin, glucagon, gastrin, and pancreatic polypeptide secretion in hemodialysis patients. *Am J Physiol*. 1994;267:F853–860. [PubMed]

- end stage renal disease. *Contrib Nephrol.* 1990;87:42–51. [PubMed]
33. Chagnac A, Weinstein T, Zevin D, et al. Effects of erythropoietin on glucose tolerance in *Clin Nephrol.* 1994;42:398–400. [PubMed]
34. Runyan JW, Hurwitz D, Robbins SL. Effect of Kimmelstiel-Wilson syndrome on insulin *N Engl J Med.* 1955;252:388–391. [PubMed]
35. Weinrauch LA, Healy RW, Leland OS Jr, et al. Decreased insulin requirements in acute nephropathy. *Arch Intern Med.* 1978;138:399–400. [PubMed]
36. Peitzman SJ, Agarwal BN. Spontaneous hypoglycemia in end stage renal failure. *Nephrol* [PubMed]
37. Arem R. Hypoglycemia associated with renal failure. *Endocrinol Metab Clin North Am.* [PubMed]
38. Ansari A, Thomas S, Goldsmith D. Assessing glycemic control in patients with diabetes failure. *Am J Kidney Dis.* 2003;41:523–531. [PubMed]
39. de Boer MJ, Miedema K, Casparie AF. Glycosylated haemoglobin in renal failure. *Diabet* 440. [PubMed]
40. Scott MG, Hoffman JW, Meltzer VN. Effects of azotemia on results of the boronate-agar exchange methods for glycated hemoglobin. *Clin Chem.* 1984;30:896–898. [PubMed]
41. Wettre S, Lundberg M. Kinetics of glycosylated hemoglobin in uremia determined on ion chromatography. *Diabetes Res.* 1986;3:107–110. [PubMed]
42. Paisley R, Banks R, Holton R, et al. Glycosylated haemoglobin in uraemia. *Diabet Med.* [PubMed]
43. Joy MS, Cefalu WT, Hogan SL, Nachman PH. Long-term glycemic control measurement receiving hemodialysis. *Am J Kidney Dis.* 2002;39:297–307. [PubMed]
44. Goldstein DE, Little RR, Lorenz RA, Malone JJ, Nathan D, Peterson CM. Test of glycemic *Care.* 2004;27(Suppl 1):S91–S93. [PubMed]
45. Position statement. Standard of Medical Care in Diabetes-2007. *Diabetes Care.* 2007;30
46. Tzamaloukas AH, Murata GH, Zager PG, Eisenberg B, Avasthi PS. The relationship between morbidity and mortality for diabetics on dialysis. *ASAIO J.* 1993;39:880–885. [PubMed]
47. Tzamaloukas AH. The use of glycosylated hemoglobin in dialysis patients. *Semin Dial.*
48. Gin H, Rigaleau V, Aparicio M. Which diet for diabetic patients with chronic renal failure *Transplant.* 1999;14:2577–2579. [PubMed]
49. Gin H, Aparicio M, Aubertin J. Low protein and low phosphorus diet in patients with chronic influence on glucose tolerance and tissue-insulin sensitivity. *Metabolism.* 1987;11:1080

- Gin H, Rigalleau V, Aparicio M. Which diet for diabetic patients with chronic renal fail
Transplant. 1999;14:2577–2579. [PubMed]
51. Snyder RW, Berns JS. Use of insulin and oral hypoglycemic medications in patients wit
advanced kidney disease. *Semin Dial.* 2004;17:365–370. [PubMed]
52. Amoff, GR.;Berns, JS.; Brier, ME. *Drug prescribing in renal failure: dosing guidelines*
Philadelphia: American College of Physicians; 1999.
53. Skillman TG, Feldman JM. The pharmacology of sulphonylureas. *Am J Med.* 1981;70:3
53. Hasslacher C. Safety and efficacy of repaglinide in type 2 diabetic patients with and wit
function. *Diabetes Care.* 2003;26:886–891. [PubMed]
54. Inoue T, Shibahara N, Miyagawa Ketal. Pharmacokinetics of nateglinide and its metabo
diabetes mellitus and renal failure. *Clin Nephrol.* 2003;60:90–95. [PubMed]
55. Nagai T, Imamura M, Iizuka K, Mori M. Hypoglycemia due to nateglinide in diabetic p
failure. *Diabetes Res Clin Pract.* 2003;59:191–194. [PubMed]
56. Charpentier G, Riveline JP, Varroud-Vial M. Management of drugs affecting blood gluc
with renal failure. *Diabetes Metab.* 2000;26(Suppl 4):73–85. K/D. [PubMed]
57. Sambanis C, Tziomalos K, Kountana E, et al. Effect of pioglitazone on heart function at
natriuretic peptide levels of patients with type 2 diabetes. *Acta Diabetol.* 2007 (In press)
59. Sampanis, Ch.,Tziomalos, K.,Kehagia, T., et al. Heart failure after thiazolidinedione the
2005;9:8–91.
60. K/DOQI clinical practice guidelines for cardiovascular disease in dialysis patients. *Am J*
3):S1–S360.
61. Tunbridge FK, Newens A, Home PD, et al. A comparison of human ultralente- and lent
injection regimens. *Diabet Med.* 1989;6:496–501. [PubMed]
62. Freeman SL, OBrien PC, Rizza RA. Use of human ultralente as the basal insulin compo
patients with IDDM. *Diabetes Res Clin Pract.* 1991;12:187–192. [PubMed]
63. Diaz-Buxo JA. Blood glucose control in diabetics: I. *Semin Dial.* 1993;6:392–397.
64. Daniels ID, Markell MS. Blood glucose control in diabetics: II. *Semin Dial.* 1993;6:394
65. Tzamaloukas AH, Orcopoulos DG. Subcutaneous versus intraperitoneal insulin in the r
CAPD: a review. *Adv Perit Dial.* 1991;7:81–85. [PubMed]
66. Selgas R, Lopez-Riva A, Alvaro F, et al. Insulin influence on the mitogenic-induced eff
effluent in CAPD patients used as an additive to dialysate. *Adv Perit Dial.* 1989;5:161–

Articles from *Hippokratia* are provided here courtesy of
Hippokratia General Hospital of Thessaloniki

[Write to PMC](#) | [PMC Home](#) | [PubMed](#)
[NCBI](#) | [U.S. National Library of Medicine](#)
[NIH](#) | [Department of Health and Human Services](#)
[Privacy Policy](#) | [Disclaimer](#) | [Freedom of information Act](#)